



Bi-Test™ CD8 FITC - CD38 PE

Product: Anti-human T and NK Lymphocytes and CD38 receptor Monoclonal Antibody.

Description: Identification of human cytotoxic/suppressor T cells expressing the 32 and 43,000 M.W. surface antigen. CD38 antigen is an integral membrane glycoprotein, M.W. 45 KDa. The CD38 antigen is present on approximately all pre-B lymphocytes, plasma cells and thymocytes. CD38 antigen is also expressed on activated T Lymphocytes, NK Cells, monocytes, myeloblasts and erythroblasts. CD38 is co-expressed on CD34+ cells, but is not expressed on pluripotent stem cells. CD38 antigen is present on T and B lymphoblastic leukemia (ALL), Burkitts' lymphoma, multiple myeloma and acute myeloid leukemia (AML).

Isotype: Mouse IgG1 kappa and IgG1 kappa.

Clones: 17D8(CD8 FITC) and TME-2 (CD38 PE).

Applications: Monitoring of activated T cell subsets in peripheral blood; Characterization of subtypes of T cell leukemias and lymphomas; Study of T lymphocyte cytokine function; Study of B cell activation; Study of AIDS virus infection; Analysis of NK cells subsets; Study of cell mediated cytotoxicity; Study of Cell-adhesion molecules relating to T & B lymphocytes.

Use: PBMC: Add 10 µl of MAB/10⁶ PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2^o to 8^oC. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature (20^oC). Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturer's instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.

Storage: Antibodies are supplied in PBS, 0.08% sodium azide and 0.2% protein carrier for FITC and PE. Antibodies should be stored at 4-8^oC. Monoclonal antibodies should not be frozen. Reagents are stable for the period shown on the vial label when stored properly.

Ordering Information:	Form	Vial Size	Catalog #
	Bi-Test™	50 Test	0838s
	Bi-Test™	100 Test	0838

For research use only. Not for use in human diagnostics or therapeutics.

REFERENCES:

1. Evolutionary Conservation of Surface Molecules that Distinguish T Lymphocyte Helper/Inducer and T Cytotoxic/Suppressor Subpopulations in Mouse and Man. Ledbetter, J.A., Evans, R.L., Lipinski, M., Cunningham-Rundles, C., Good, R.A. and Herzenberg, L.A., J. Exp. Med. 153,310, 1981.
2. Circulating Antigen-Specific Suppressor T Cells in a Healthy Woman: Mechanism of Action and Isolation with a Monoclonal Antibody. Engleman, E.G., Benike, C.J., and Evans, R.L., Clin. Res. 29, 365a 1981.
3. Induction of Immunoglobulin Secreting Cells in the Allogeneic Mixed Leukocyte Reaction: Regulation by Helper and Suppressor Lymphocyte Subsets in Man. Kotzin, B.L., Benike, C.J. and Engleman, E.G., J. Imm. 127,931,1981.
4. CD4 and CD8 molecules can physically associate with the same T-cell receptor. Gallagher, P.F., Fazekas de St. Groth, B., Miller, J.F.A.P., Proc. Nat. Acad. Sci. 1989,86:10044.
5. Immunofluorescence Measurement in a Flow Cytometer using Low-Power Helium Neon Laser Excitation. Shapiro, H.M, Glazer, A.N., Christenson, L., Williams, J.M., and Strom, T. B. Cytometry 4,276, 1983.
6. Comparison of Helium Neon and Dye lasers for Excitation of Allophycocyanin. Loken, M.R., Kiej, J.F. and Kelly, K.,A. Cytometry 8, 96, 1987.
- 7) CD38 Molecule: Structural and Biochemical Analysis on Human T Lymphocytes, Thymocytes and Plasma Cells. J. Immunol, 1990 Aug; 145(3):878-84. Alessio, M. Roggero, S.; Funaro, A. ; De Monte L.B.; Peruzzi, L.; Geuna, M.; Malavasi, F.
- 8) Differences Among Mononuclear Cell Subpopulations in HIV Seropositive or Seronegative Homosexual and Heterosexual men as Determined by Four Color Flow Cytometry. J. Acquir Immune Defic Syndr., 1990;3(8):747-56. Schnizlein-Bick, C.T.; Magier, M.R.; Jones, R.B.; Fife, K.H.; Katz, B., P.; Walker, E.B.
- 9) Quantitative Comparison of Myeloid Antigens on Five Lineages of Mature Peripheral Blood Cells. J. Leukoc Bio. 1990, Au;48(2):138-48. Terstappen, L.W.; Hollander, Z.; Meiners, H.; Loken, M.R.
- 10) B-cell Surface Phenotypes of Proliferating Myeloma Cells: Target Antigens for Immunotherapy. Am. J. Hematol. 1990, Fe;33(2):101-9, Chan, C.S; Wormsley, S.B.; Pierce, L. E.; Peter, J.B.; Schechter, G.P.
- 11) Interrelationships Between Serologic Markers of Immune Activation and T Lymphocyte Subsets in HIV Infection. J. Acquir Immune Defic Syndr. 1990;3(5):525-30. Prince, H.E.; Kleinman, S. Czaplicki, C.; John, J.; Williams, A.E.
- 12) Sequential Generations of Hematopoietic Colonies Derived from Single Nonlineage-committed CD34+ CD38- Progenitor Cells. Blood 1991, Mar 1;77(6):1218-27. Terstappen, L.W.; huang, S.; Safford, M. Lansdorp, P.M.; Loken, M.R.
- 13) Involvement of Multilineage CD38 Molecule in a Unique Pathway of Cell Activation and Proliferation. J. Immunol, 1990 Oct 1;145(8):2390-6Funaro, A.; Spagnoli, G.C.; Ausiello, C.M.; Alessio, M.; Roggero, S.; Delia, D.; Zaccolo, M.; Malavasi, F.
- 14) Induction of CD4 and Susceptibility to HIV-1 Infection in Human CD8+ T Lymphocytes by Human Herpesvirus 6. Nat.349(6309): 533-5, 1991. Lusso, P.; De Maria, A.; Malnati, M.; Lori, F.; DeRocco, S.E.;Baseler, M.; Gallo, R.C.
- 15) Evolutionary Conservation of Surface Molecules That Distinguish T Lymphocyte Helper/Inducer and T Cytotoxic/Suppressor Subpopulations in Mouse and Man. J. Exp Med 153,310, 1981. Ledbetter, J. A.; Evans, R.L.; Lipinski, M.; Cunningham-Rundles, C.; Good, R.A; Herzenberg, L. A.
- 16) Circulating Antigen-Specific Suppressor T Cells in a Healthy Woman: Mechanism of Action and Isolation with a Monoclonal Antibody. Clin. Res. 29, 365a, 1981. Engleman, E. G.; Benike, C. J.; Evans, R.L
- 17) Induction of Immunoglobulin Secreting Cells in the Allogeneic Mixed Leukocyte Reaction: Regulation by Helper and Suppressor Lymphocyte Subsets in Man. J. Imm.127,931, 1981. Kotzin, B.L.; Benike, C.J.; Engleman, E. G.
- 18) CD4 and CD8 Molecules Can Physically Associate with the Same T-cell Receptor. Proc. Nat. Acad. Sci. 1989, 86:10044.

For research use only. Not for use in human diagnostics or therapeutics.

Exalpha Biologicals, Inc., 86 Rosedale Rd. Watertown, MA 02472
Tel: 800.395.1137 or 617.924.3400, Fax: 866.924.5100 or 617.924.5100, Web:www.exalpha.com