



Bi-Test™ CD4 FITC - CD45RA PE

Product: Anti-Human CD4 (FITC) Helper/Inducer T Cell Monoclonal Antibody and Anti-human CD45RA (PE) Leucocyte Common Antigen Monoclonal Antibody.

Description: Identification of CD4 human helper/inducer T cells expressing the 60,000 M.W. surface antigen (HLA class II reactive). CD4 is present in low density on monocytes. CD45RA recognizes Mr. 220-kdalton of the leucocyte antigen (LCA). CD45RA antigen is a member of the CD45 antigen family that also includes CD45, CD45RO, and CD45RB antigens. CD45RA is Expressed on approximately 75% of CD8 lymphocytes and 50% of CD4 lymphocytes, all B cells and NK Cells. CD45RA antigen is expressed on naive T lymphocytes, antigen density decreases upon in vitro activation. A decrease antigen density has been demonstrated for CD4+CD45RA+subset with active multiple sclerosis.

Isotype: Mouse IgG1 kappa (FITC) and Mouse IgG1 kappa (PE)

Clones: 7E14 (CD4 FITC) and BHKG-2 (CD45RA PE)

Applications: Monitoring of T cells subsets in peripheral blood; Characterization of subtypes of T cell leukemias and lymphomas; Analysis of T cell subsets involved in helper/inducer functions; Studies of AIDS virus infection; Study of T lymphocyte cytokine function; Study of B cell activation; Study of Cell-adhesion molecules relating T and B lymphocytes; Studies of multiple sclerosis and other autoimmune disease.

Use: PBMC: Add 10 μ l of MAB/10⁶ PBMC in 100 μ l PBS. Mix gently and incubate for 15 minutes at 20^o to 8^oC. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 μ l of MAB/100 μ l of whole blood. Mix gently and incubate for 15 minutes at room temperature 20^oC. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturer's instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.

Storage: Antibodies are supplied in PBS, 0.08% sodium azide and 0.2% protein carrier for FITC and PE. Antibodies should be stored at 4-8^oC. Monoclonal antibodies should not be frozen. Reagents are stable for the period shown on the vial label when stored properly.

Ordering Information:	Form	Vial Size	Catalog #
	Bi-Test™	50 Test	445RAs
	Bi-Test™	100 Test	445RA

For research use only. Not for use in human diagnostics or therapeutics.

REFERENCES:

1. Thymus Dependent Membrane Antigens in Man: Inhibition of Cell-Mediated Lympholysis by Monoclonal Antibodies to the TH-2 Antigen. Evans,R.L., Wall,D.W., Platsoucas,C.D., Siegal,F.P., Fikrig,S.M., Testa,C.M, and Good,R.A. Proc. Nat. Acad. Sci. 78,544,1981
2. Novel Immunoregulatory Functions of Phenotypically Distinct Subpopulations of CD4+ cells in the Human Neonate. Clement,L.T., Vink,P.E., Bradley,G.E. J. Immunology 145(1):102-8,1990
3. Antigen Presentation by the CD4 Positive Monocyte Subset. Szabo,G., Miller,C.L., Kodys,K., J. Leukoc. Biol. 47(2): 111-20,1990
4. Human Immunodeficiency Virus Infection is Efficiently Mediated by a Glycolipid-Anchored form of CD4. Diamond,D.C., Finberg,R., Chaudhuri,S., Sleckman,B.P., Burakoff,S.J., Proc. Natl. Acad. Sci. 87(13):5001-5,1990
5. Development Regulation of the Intrathymic T cell Precursor Population. Adkins,B., J. Immunol. 146(5):1387-93,1991
6. Leucocyte TypingIV: White Cell Differentiation Antigens, Schwinzer R. Cluster report: CD45/CD45R, Knapp W., Dorken B., Gilks W.R.,et al, eds. Oxford University Press; 1989:628-634
7. CD45 isoform expression on human neonatal T cells: expression and turnover of CD45 isoforms on neonatal versus adult T cells after activation. Cell Immunol 1992 Jun;142(1):114-24 Yamada A; Kaneyuki T; Hara A; Rothstein DM; Yokoyama MM
8. Transient accumulation and subsequent rapid loss of messenger RNA encoding high molecular mass CD45 isoforms after T cell activation. J Immunol 1992 Mar 15;148(6):1898-905 Deans JP; Serra HM; Shaw J; Shen YJ; Torres RM; Pilarski LM
9. Patterns of membrane CD45 isoform expression by leukaemic blasts and normal mature myeloid cells. Int J Hematol 1992 Jun;55(3):235-42 Master PS; Richards SJ; Roberts BE; Scott CS
10. CD45 isoform expression during T cell development in the thymus Eur J Immunol 1992 Jul;22(7):1843-50 Fujii Y; Okumura M; Inada K; Nakahara K; Matsuda H
11. Functional subsets of T cells defined by isoforms of CD45 Biochem Soc Trans 1992 Feb;20(1):184-7 Beverley PC; Daser A; Michie CA; Wallace DL

For research use only. Not for use in human diagnostics or therapeutics.

Exalpha Biologicals, Inc., 86 Rosedale Rd. Watertown, MA 02472
Tel: 800.395.1137 or 617.924.3400, Fax: 866.924.5100 or 617.924.5100, Web:www.exalpha.com