



Spingosine 1 Phosphate 4 Receptor (CT) (EDG6) Blocking Peptide

Blocking peptide for use in Western blotting with Anti-EDG6 (S1P4) (CT) (Cat. No. X1182P)

CAT. No.
X1312B
Size
50 µg

CONCENTRATION

1 mg/ml

APPLICATIONS

Antibody function blocking

COMMENTS

For use with EDG6 (S1P4) (CT) polyclonal antibody (Cat. No. X1182P).

FORMULATION

Provided in solution at 1 mg/ml in phosphate buffered saline.

STORAGE CUSTOMER

Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

USE

Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37 deg C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 µl or 10 µg of Exalpha's rabbit anti-EDG6 (S1P4) (CT) (Cat. No. X1182P) is added to 10 µg of blocking peptide for a total volume of 20 µl. The mixture is allowed to incubate for 20 minutes at 37 deg C prior to dilution in suitable buffer (for Western blot, etc.).