

pS2 Estrogen-Regulated Protein. Mouse, Clone pS2.1 Monoclonal Antibody, Human

BACKGROUND

It recognizes a polypeptide of 6.5kDa which is identified as pS2 estrogen-regulated protein. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy.

ORDERING INFORMATION

CATALOG NUMBER
X1440M

SIZE
100 µg

FORM
Purified

HOST/CLONE
Mouse, Clone pS2.1

FORMULATION
Provided as solution in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION
1 mg/ml

ISOTYPE
IgG1

APPLICATIONS
Frozen Section Staining
Paraffin Section Staining

IMMUNOGEN


Hybridoma produced by the fusion of splenocytes from BALB/c mice immunized with synthetic peptide of 31 amino acids from the C-terminus of human pS2 protein and mouse myeloma p3-NS1-Ag4-1 cells.

SPECIES REACTIVITY

Human

Legend:

Immunohistochemical staining using pS2 Estrogen-Regulated Protein antibody on formalin fixed, paraffin embedded human breast carcinoma.



For research use only. Not for use in human diagnostics or therapeutics.

Exalpha Biologicals, Inc. 86 Rosedale Road Watertown, MA 02472
Tel: 800.395.1137 Fax: 866.924.5100 www.exalpha.com info@exalpha.com

Page 1 of 2
Cat. No. X1440M

POSITIVE CONTROL

Breast carcinomas

COMMENTS

This antibody can be used for immunohistochemistry on frozen and formalin/paraffin embedded tissues (2-4 µg/ml). Optimal concentration should be evaluated by serial dilutions.

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -20 degrees C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

REFERENCES

1. Masaiakowski P, et al. Cloning of cDNA sequences of hormone-regulated genes from the MCF-7 human breast cancer cell line. *Nucleic Acids Res* 10:7895-7903, 1982.
2. Jeltsh JM, et al. Structure of human oestrogen-responsive gene pS2. *Nucleic Acids Res* 15:1401-1414, 1987.
3. Mori K, et al. Complete primary structure of the oestrogen-responsive gene (pS2) product. *J Biochem* 107:73-76, 1990
4. Rio MC, et al. Breast cancer-associated pS2 protein: synthesis and secretion by normal stomach mucosa. *Science* 241:705-708, 1988.
5. Rio MC, et al. Specific expression of the pS2 gene in subclasses of breast cancers in comparison with expression of the estrogen and progesterone receptors and the oncogene ERB B2, *Proc Natl Acad Sci USA* 84:9243-9247, 1987.
6. Molina R, et al. Estrogen-inducible pS2 protein: Tumor characteristics and clinical behavior in breast cancer. *Proc Endocrine Soc* 61(#148), 1990.
7. Abbondanzo SL, et al. Prognostic significance of immuno-cytochemically determined pS2 in axillary node-negative breast carcinoma. *Breast Cancer Res Treat* 16:182, 1990.

LAST MODIFIED 9/16/2002