

Interleukin-2. Rat Monoclonal Antibody JES6-5H4 Mouse

IL-2

BACKGROUND

Interleukin-2 (IL-2) is produced by T-cells in response to antigenic or mitogenic stimulation. This protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. IL-2 can stimulate B-cells, monocytes, lymphokine-activated killer cells, natural killer cells, and glioma cells.

ORDERING INFORMATION

CATALOG NUMBER
X1924M

SIZE
0.5 mg

FORM
Unconjugated

HOST/CLONE
Rat Clone JES6-5H4

FORMULATION
Purified IgG in phosphate buffered saline, pH 7.4

CONCENTRATION
1 mg/ml

ISOTYPE
IgG2b

APPLICATIONS
ELISA, Immunoprecipitation, Flow Cytometry

IMMUNOGEN

Hybridoma produced by the fusion of splenocytes from rat immunized with mouse recombinant IL-2 protein and P3X63-AG8.653 myeloma cells.

SPECIES REACTIVITY

Mouse

COMMENTS

Antibody can be used for ELISA as the detector antibody, immunoprecipitation and flow cytometry. Antibody is non-neutralizing. Optimal concentration should be evaluated by serial dilutions.

STORAGE

Product should be stored at -70°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

For research use only. Not for use in human diagnostics or therapeutics.

POSITIVE CONTROL/TISSUE EXPRESSION

Detects natural and recombinant mouse IL-2

SHIP CONDITIONS

Ship on dry ice, freeze upon arrival

REFERENCES

Abrams, J.S., M.G. Roncarolo, H.Yssel, U. Andersson, G.J. Gleich, and J.E. Silver (1992) Strategies of anti-cytokine monoclonal antibody development. Immunological Review 127:5–24.

LAST MODIFIED 2/19/2008

For research use only. Not for use in human diagnostics or therapeutics.

Exalpha Biologicals, Inc. 5 Clock Tower Place, Suite 255 Maynard, MA 01754
Tel: 800.395.1137 Fax: 866.924.5100 www.exalpha.com info@exalpha.com

Page 2 of 2
Cat. No. X1924M