



## EDG8 (S1P5) Control Lysate

### BACKGROUND

Endothelial Cell Differentiation Gene-8 (EDG8 (S1P5)) belongs to a family of G-protein coupled receptors whose ligands are lysophospholipids. The ligand for EDG8 (S1P5) is sphingosine-1-phosphate. There are 8 known members of the EDG receptor family and they are implicated in mediating growth related effects such as induction of cellular proliferation, alterations in differentiation and survival and suppression of apoptosis. They also evoke cellular effector functions that are dependent on cytoskeletal responses such as contraction, secretion, adhesion and chemotaxis. EDG receptors are developmentally regulated and differ in tissue distribution. They couple to multiple types of G proteins to signal through ras and MAP kinase, rho, phospholipase C and several protein tyrosine kinases. EDG8 (S1P5) is expressed in oligodendrocytes and fibrous astrocytes in the rat brain.

### ORDERING INFORMATION

**CATALOG NUMBER**  
X1487C

**SIZE**  
100  $\mu$ l

**CUSTOMER STORAGE**  
Product should be stored at -20°C.  
Aliquot to avoid freeze/thaw cycles

**FORMULATION**  
Provided in 10% glycerol, 0.063 M Tris-HCl (pH 6.8), 2% SDS and 0.002% bromophenol blue, 5% 2-mercaptoethanol

**SHIP CONDITIONS**  
Ship at ambient temperature, freeze upon arrival

**STABILITY**  
Products are stable for one year from purchase when stored properly

### COMMENTS

Optimal concentration should be evaluated by serial dilutions.

### INSTRUCTIONS

Use 10-20  $\mu$ l of EDG8 (S1P5) cell lysate (Cat. No. X1487C) per lane as a control for using EDG8 (S1P5) antibody (Cat. No. X1094P). Due to low expression of EDG receptors, we recommend use of Pierce Femto Signal substrate for western blot development.