



MEG-2 (285-593)/PTPN9. E.coli Active Enzyme Protein-tyrosine phosphatase MEG2, PTPase-MEG2

BACKGROUND

MEG-2 also known as PTPN9 is a protein-tyrosine phosphatase originally isolated from peripheral neutrophilic polymorphonuclear leukocytes (PMN). MEG2 is predominantly cytosolic with components present in secondary and tertiary granules and secretory vesicles. MEG2 activity is inhibited after exposure of cells to opsonized zymosan or to phorbol 12-myristate 13-acetate but largely unaffected by the f-met-leu-phe (N-formyl-methionyl-leucyl-phenylalanine). Cysteine 515 is essential for catalytic activity of MEG-2, whereas the noncatalytic (N-terminal) domain of MEG2 negatively regulates the enzymatic activity of the C-terminal phosphatase domain. The activity of MEG2 is enhanced by specific polyphosphoinositides. MEG2 associates at an early stage with nascent phagosomes. MEG2 is a polyphosphoinositide-activated tyrosine phosphatase that may be involved in signaling events regulating phagocytosis, and essential antimicrobial function in the immune response.

ACTIVITY

1070 pmole/min/ μ g of enzyme; Determined using pNPP; Reaction conditions: 1 mM pNPP, 15 min incubation at room temperature, 2 ng/ μ l enzyme.

PURITY

>90% pure as determined by Coomassie-stained SDS gel

APPLICATIONS

ORDERING INFORMATION

CATALOG NUMBER

X1662E

SIZE

20 μ g

CUSTOMER STORAGE

Enzyme should be stored at -20°C.
Enzyme should be kept on ice when dispensing

FORMULATION

Provided in 25 mM Tris-HCl, pH 8.0,
67 mM NaCl, 50% glycerol, 1.4 mM
KCl, 3 mM DTT, 10 mM glutathione.

SHIP CONDITIONS

Ship on gel ice, store at -20°C
immediately upon arrival

STABILITY

Products are stable for one year from
purchase when stored properly

CONCENTRATION

See vial for concentration

SOURCE

Recombinant enzyme produced in E.
coli

ASSAY METHODS

MATERIALS

1. Assay Buffer: 50 mM HEPES, pH 7.4, 100 mM NaCl, 2 mM EDTA, 3 mM DTT
2. Stop solution: 2M K₂CO₃
3. 190 mM pNPP
4. Microtiter plate
5. Microtiter plate reader capable of measurements at 405 nm
6. Water bath or incubator at 30°C

PROCEDURE

1. Prepare reaction mixture:
 - a. 73 μ l assay buffer
 - b. 26 μ l pNPP (Final concentration of pNPP is 50 mM)
 - c. 1 μ l of MEG-2
2. Mix well and start reaction at 30°C in water bath and incubate for 10 min.
3. Add 100 μ l per well of 2 M K₂CO₃ to stop the reaction.
4. Read absorbance at 405 nm using a microtiter plate reader.

REFERENCES

- 1: Huynh H, Bottini N, Williams S, Cherepanov V, Musumeci L, Saito K, Bruckner S, Vachon E, Wang X, Kruger J, Chow CW, Pellecchia M, Monosov E, Greer PA, Trimble W, Downey GP, Mustelin T. Control of vesicle fusion by a tyrosine phosphatase. *Nat Cell Biol.* 2004 Sep;6(9):831-9. Epub 2004 Aug 22.
- 2: Huynh H, Wang X, Li W, Bottini N, Williams S, Nika K, Ishihara H, Godzik A, Mustelin T. Homotypic secretory vesicle fusion induced by the protein tyrosine phosphatase MEG2 depends on polyphosphoinositides in T cells. *J Immunol.* 2003 Dec 15;171(12):6661-71.
- 3: Xu MJ, Sui X, Zhao R, Dai C, Krantz SB, Zhao ZJ. PTP-MEG2 is activated in polycythemia vera erythroid progenitor cells and is required for growth and expansion of erythroid cells. *Blood.* 2003 Dec 15;102(13):4354-60. Epub 2003 Aug 14.
- 4: Zhao R, Fu X, Li Q, Krantz SB, Zhao ZJ. Specific interaction of protein tyrosine phosphatase-MEG2 with phosphatidylserine. *J Biol Chem.* 2003 Jun 20;278(25):22609-14. Epub 2003 Apr 17.
- 5: Qi Y, Zhao R, Cao H, Sui X, Krantz SB, Zhao ZJ. Purification and characterization of protein tyrosine phosphatase PTP-MEG2. *J Cell Biochem.* 2002;86(1):79-89.
- 6: Wang X, Huynh H, Gyorloff-Wingren A, Monosov E, Stridsberg M, Fukuda M, Mustelin T. Enlargement of secretory vesicles by protein tyrosine phosphatase PTP-MEG2 in rat basophilic leukemia mast cells and Jurkat T cells. *J Immunol.* 2002 May 1;168(9):4612-9.
- 7: Kruger JM, Fukushima T, Cherepanov V, Borregaard N, Loeve C, Shek C, Sharma K, Tanswell AK, Chow CW, Downey GP. Protein-tyrosine phosphatase MEG2 is expressed by human neutrophils. Localization to the phagosome and activation by polyphosphoinositides. *J Biol Chem.* 2002 Jan 25;277(4):2620-8. Epub 2001 Nov 15.