



Sirtuin 1 (193-741). E.coli Active Enzyme HDAC Type 3,-1; Silent mating type information regulation 2 homolog 1

BACKGROUND

Human Sirtuin 1 (GenBank Accession No. NM_012238), amino acids 193-741 with N-terminal GST tag, MW=87.2 kDa, expressed in E.coli expression system.

Sirtuin 1 is a NAD-dependent deacetylase, which regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. Deacetylates Lys-382 of p53/TP53 and impairs its ability to induce proapoptotic program and modulate cell senescence. Deacetylates TAF1B and thereby represses rDNA transcription by the RNA polymerase I. Involved in HES1- and HEY2-mediated transcriptional repression. Inhibits skeletal muscle differentiation by deacetylating PCAF and MYOD1. May serve as a sensor of the cytosolic ratio of NAD(+)/NADH, which is essential in skeletal muscle cell differentiation. Despite some ability to deacetylate histones in vitro, such activity is either weak or nonexistent in vivo. Inhibited by nicotinamide. Activated by resveratrol (3,5,4'-trihydroxy-trans-stilbene), butein (3,4,2',4'- tetrahydroxychalcone), piceatannol (3,5,3',4'-tetrahydroxy-trans-stilbene), Isoliquiritigenin(4,2',4'-trihydroxychalcone), fisetin (3,7,3',4'-tetrahydroxyflavone) and quercetin (3,5,7,3',4'- pentahydroxyflavone).³

ACTIVITY

Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling. 10.2 pmol/min/ μ g. One U =1 pmol/min, Assay condition: 25 mM Tris/HCl, pH 8.0, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl₂, and 0.1 mg/ml BSA, 20 μ M HDAC substrate 1, 500 μ M NAD⁺ and Sirtuin 1. Incubation condition: 30 min at 37°C followed by

PURITY

> 60%

APPLICATIONS

ORDERING INFORMATION

CATALOG NUMBER
X2037E

SIZE
100 μ g

CUSTOMER STORAGE
Product should be stored at -80°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION
Provided in 50 mM Tris-HCl, pH 8.0,
138 mM NaCl, 20 mM glutathione, and
10% glycerol.

SHIP CONDITIONS
Ship on dry ice, freeze upon arrival

STABILITY
Stable for > 6 months when stored as
recommended.

CONCENTRATION
Lot Specific

SOURCE

ASSAY METHODS

MATERIALS

1. 4 ul of Sirtuin 1 (0.25 ug/ul)
2. 31 ul of HDAC assay buffer (Exalpha catalog number xxxx)
3. 5 ul of 1 mg/ml BSA
4. 5 ul of 2 mM NAD+
5. 5 ul of 300 uM substrate (Biomol catalog number: KI-177 gave the best results)

PROCEDURE

1. Add all reaction mixture to a low binding black plate
2. Incubate at 30°C for 60 minutes
3. Stop the reaction by adding 50 ul of HDAC assay developer and incubate the plate at rt for 15 minutes.
4. Read sample in microplate reader at a wavelength of 350-380 nm and detection of emitted light at 440-460nm.

REFERENCES

1. Kolthur-Seetharam, U. et al (2006) Cell Cycle 5, 873-877.
2. Solomon, J.M. et al(2006) Mol. Cell. Biol. 26, 28-38.
3. UCSC Genome Browser-<http://genome.ucsc.edu/>., Specific for Sirtuin 1:
http://genome.ucsc.edu/cgi-bin/hgGen?db=hg18&hgg_gene=NM_012238&hgg_chrom=chr10&hgg_start=69314432&hgg_end=69348147, Accessed March 13, 2007.

PRODUCT SPECIFIC REFERENCES

Last Modified
5/30/2017

Page 2 of 2
Cat. No. X2037E